


Allergy Action Plan

Must be accompanied by a Medication Authorization Form (OCC 1216)



TREATMENT

CHILD'S NAME: _____ Date of Birth: _____

ALLERGY TO: _____

Is the child Asthmatic? No Yes (If Yes = Higher Risk for Severe Reaction)

Symptoms:	Give this Medication	
The child has ingested a food allergen or exposed to an allergy trigger:	Epinephrine	Antihistamine
But is not exhibiting or complaining of any symptoms		
Mouth: itching, tingling, swelling of lips, tongue or mouth		
Skin: hives, itchy rash, swelling of the face or extremities		
Gut: nausea, abdominal cramps, vomiting, diarrhea		
Throat*: tightening of throat, hoarseness, hacking cough		
Lung*: shortness of breath, repetitive coughing, wheezing		
Heart*: fainting, pale, blueness		
Other:		
If reaction is progressing (several of the above areas affected)		

*Potentially life-threatening. The severity of symptoms can quickly change.

Medication	Dose:
Epinephrine:	
Antihistamine:	
Other:	

IMPORTANT: Asthma inhalers and/or antihistamines cannot be depended on to replace epinephrine in anaphylaxis.

Doctor's Signature

Date

EMERGENCY CALLS

1) Call 911 (or Rescue Squad) whenever Epinephrine has been administered. 2) Call the parent. State that an allergic reaction has been treated and additional epinephrine may be needed.

Doctor's Name: _____

Phone Number: _____

Contact(s)	Name/Relationship	Phone Number(s)	
		Daytime Number	Cell
Parent/Guardian 1			
Parent/Guardian 2			
Emergency 1			
Emergency 2			

EVEN IF A PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE AND CALL 911.

Parent/Guardian's Signature

Date

Allergy Action Plan (Continued)

Must be accompanied by a Medication Authorization Form (OCC 1216)



CHILD'S NAME: _____ Date of Birth: _____

ALLERGY TO: _____

Is the child Asthmatic? No Yes (If Yes = Higher Risk for Severe Reaction)

The Child Care Facility will:

- Reduce exposure to allergen(s) by: (no sharing food,
- Ensure proper hand washing procedures are followed.
- Observe and monitor child for any signs of allergic reaction(s).
- Ensure that medication is immediately available to administer in case of an allergic reaction (in the classroom, playground, field trips, etc.)
- Ensure that a person trained in Medication Administration accompanies child on any off-site activity.
-

EPIPEN®
(Epinephrine) Auto-Injectors 0.3/0.15mg

userguide

1 Pull off the blue safety release cap.

2 Swing and firmly push the orange tip against the outer thigh so it 'clicks.' HOLD on thigh for approximately 10 seconds to deliver the drug.
Please note: As soon as you release pressure from the thigh, the protective cover will extend.
Each EpiPen Auto-Injector contains a single dose of a medicine called epinephrine, which you inject into your outer thigh. DO NOT INJECT INTRAVENOUSLY. DO NOT INJECT INTO YOUR BUTTOCK, as this may not be effective for a severe allergic reaction. In case of accidental injection, please seek immediate medical treatment.

Call 911

3 Seek immediate emergency medical attention and be sure to take the EpiPen Auto-Injector with you to the emergency room.

To view an instructional video demonstrating how to use an EpiPen Auto-Injector, please visit epipen.com.

©2010 Dey Pharma, L.P. All rights reserved.
 DCV® and the Dey logo are registered trademarks of Dey Pharma, L.P.
 EpiPen®, Epi-Pen 2-Pak®, and Epi-Pen Jr 2-Pak® are registered trademarks of Mylan Inc licensed exclusively to its wholly-owned subsidiary, Dey Pharma, L.P.

The Parent/Guardian will:

- Ensure the child care facility has a sufficient supply of emergency medication.
- Replace medication prior to the expiration date
- Monitor any foods served by the child care facility, make substitutions or arrangements with the facility, if needed.
-